



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

The impact of dawn to sunset fasting on immune system and its clinical significance in Covid-19 pandemic

Sundus I. Bhatti, Ayse L. Mindikoglu



PII: S2589-9368(21)00086-4

DOI: <https://doi.org/10.1016/j.metop.2021.100162>

Reference: METOP 100162

To appear in: *Metabolism Open*

Received Date: 13 November 2021

Revised Date: 23 December 2021

Accepted Date: 24 December 2021

Please cite this article as: Bhatti SI, Mindikoglu AL, The impact of dawn to sunset fasting on immune system and its clinical significance in Covid-19 pandemic, *Metabolism Open* (2022), doi: <https://doi.org/10.1016/j.metop.2021.100162>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Inc.

## **AUTHOR CONTRIBUTIONS**

**Sundus I. Bhatti, MD**, contributed to the drafting and writing the manuscript and performed a critical review of the manuscript for important intellectual content.

**Ayse L. Mindikoglu, MD, MPH**, contributed to the drafting and writing the manuscript and performed a critical review of the manuscript for important intellectual content.

# THE IMPACT OF DAWN TO SUNSET FASTING ON IMMUNE SYSTEM AND ITS CLINICAL SIGNIFICANCE IN COVID-19 PANDEMIC

Sundus I. Bhatti, M.D.<sup>1,2</sup>, Ayse L. Mindikoglu, M.D., M.P.H.<sup>1,2</sup>

## **Institutions:**

1. Margaret M. and Albert B. Alkek Department of Medicine, Section of Gastroenterology and Hepatology, Baylor College of Medicine, Houston, TX
2. Michael E. DeBakey Department of Surgery, Division of Abdominal Transplantation, Baylor College of Medicine, Houston, TX

## **Correspondence:**

Ayse L. Mindikoglu, M.D., M.P.H.

Associate Professor of Medicine

Baylor College of Medicine

Margaret M. and Albert B. Alkek Department of Medicine

Section of Gastroenterology and Hepatology

Michael E. DeBakey Department of Surgery

Division of Abdominal Transplantation

6620 Main Street- Suite 1450

Houston, TX 77030

Phone: 832-355-1400; Fax: 713-610-2479

E-mail: Ayse.Mindikoglu@bcm.edu

**ABSTRACT**

Dawn to sunset fasting, a type of intermittent fasting commonly practiced in the month of Ramadan, requires fasting from dawn to sunset without food or liquid intake. Dawn and dusk are two transition time zones of the day that play a critical role in the human circadian rhythm. Practicing dawn to sunset fasting requires the alignment of mealtimes and wake-sleep times with the human biological dawn and dusk.

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) impairs immune cell responses at multiple levels and leads to severe Coronavirus Disease 2019 (COVID-19). It generates high levels of pro-inflammatory cytokines and chemokines, also known as a cytokine storm, leads to mitochondrial dysfunction and generation of excessive amounts of mitochondrial reactive oxygen species, downregulates autophagy to escape detection for unchecked replication, and alters gut microbiome composition. Severe cases of COVID-19 have been associated with several comorbidities that impair immune responses (e.g., obesity, diabetes, malignancy) and blood laboratory abnormalities (e.g., elevated procalcitonin, C-reactive protein, interleukin-6 (IL-6), leukocytosis, lymphopenia).

Several studies of dawn to sunset fasting showed anti-inflammatory effect by suppressing several pro-inflammatory cytokines, reducing oxidative stress, inducing a proteome response associated with increased autophagy, remodeling the gut microbiome, and improving the components of metabolic syndrome (e.g., obesity, blood glucose levels, blood pressure, lipids).

In conclusion, dawn to sunset fasting has the potential to optimize the immune system function against SARS-CoV-2 during the COVID-19 pandemic as it suppresses chronic

inflammation and oxidative stress, improves metabolic profile, and remodels the gut microbiome. This review presents scientific literature related to the effects of dawn to sunset fasting on the immune system. Studies are needed to assess and confirm the potential benefits of dawn to sunset fasting against SARS-CoV-2.

**Keywords:** COVID-19, SARS-CoV-2, coronavirus, intermittent fasting, Ramadan fasting, Ramadan-associated intermittent fasting, diurnal fasting, immune system, dawn to sunset fasting, fasting, inflammation, oxidative stress, autophagy

## 1. INTRODUCTION

According to World Health Organization (WHO), as of December 21, 2021, the Coronavirus Disease 2019 (COVID-19) pandemic has a death toll of over 5 million (1). Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has raised great interest in human immunology. There is a lot that remains unknown about SARS-CoV-2. More importantly, there is an urgency towards understanding the lifestyle and pharmacologic measures to help prevent and decrease the acuity of the infection. As we learn more about the virus, several comorbidities (e.g., advanced age, male sex, obesity, diabetes, hypertension, malignancy, thromboembolism, chronic obstructive pulmonary disease, interstitial lung disease, chronic liver disease, acute kidney injury, and chronic kidney disease) and blood laboratory abnormalities (e.g., elevated procalcitonin, basal urea nitrogen, creatinine, C-reactive protein, interleukin-6, cardiac troponin I, leukocytosis, lymphopenia, thrombocytopenia, ferritin, D-dimer levels) have been associated with increased risk of severe COVID-19 infection (2-5).

Despite the roll-out of vaccines, there is no end in sight as the virus mutates and increases virulence. A spectrum of therapeutic drugs has evolved since the beginning of the pandemic, with current treatments including, but are not limited to, antivirals, immunomodulators (e.g., corticosteroids, biologics), anticoagulants, and complement inhibitors (6). Thus far, preventative measures include social distancing, wearing proper-fitting medical masks, eye protection, frequent hand washing, and vaccination (7, 8). In addition to these preventative measures, intermittent fasting and time-restricted eating were proposed as a potential lifestyle modification to strengthen the immune system during the COVID-19 pandemic (3, 9-14). Although there are no randomized controlled studies conducted to assess the effects of intermittent fasting and time-restricted eating on the immune system during the COVID-19 pandemic, there has since been an avid interest in

the physiology of intermittent fasting and time-restricted eating as a means of modulating the immune system to viral infections, including SARS-CoV-2 (3, 9-14).

## **2. DAWN TO SUNSET FASTING AND ITS DIFFERENCE FROM OTHER TYPES OF INTERMITTENT FASTING**

Two major types of intermittent fasting have been adopted to pursue a healthier lifestyle based on the starting and ending times of the fasting period. The most popular type of intermittent fasting is dawn to sunset/dawn to dusk fasting that has one fasting and one eating/drinking period within a day. Dawn to sunset fasting is a type of intermittent fasting commonly practiced in the month of Ramadan (15). Although the term "time-restricted eating" can be used interchangeably with "intermittent fasting," time-restricted eating has a broader meaning. In general, while intermittent fasting has one fasting and one eating/drinking period within a day, time-restricted eating may have more than one fasting and one eating/drinking period within a day (e.g., breakfast between 7 am and 8 am followed by fasting from 8 am until 1 pm, lunch between 1 pm and 2 pm followed by fasting from 2 pm until 7 pm, and dinner between 7 pm and 8 pm followed by fasting from 8 pm until 7 am next day).

In dawn to sunset fasting, the daily fast starts at dawn after having a pre-dawn breakfast and ends at dusk with dinner (16, 17). Dawn and dusk are two transition time zones of the day and play a critical role in the function of the human circadian pacemaker (18, 19). Morning and evening oscillators, the two components of the circadian pacemaker, were shown to be entrained to dawn and dusk, respectively (18, 19). Therefore, starting the fast at dawn and ending it at sunset/dusk appears to be compliant with the function of these two components of the circadian pacemaker and act as a robust Zeitgeber (a time giver) (16), entraining peripheral oscillators to dawn and dusk.



Mealtimes during intermittent fasting play a critical role in maintaining a healthy circadian rhythm. During dawn to sunset fasting, mealtimes should be scheduled at the transition zones of the day, including a pre-dawn breakfast just before dawn and dinner at sunset. Pre-dawn breakfast should not be skipped as skipping breakfast was shown to cause a disrupted cortisol circadian rhythm and elevate blood pressure (20). Skipping breakfast or delaying it to a later hour in the morning during intermittent fasting can exert additional cardiometabolic stress, specifically in individuals with metabolic syndrome (20). Likewise, eating meals or having snacks in the middle of the night or throughout the night would likely disrupt the circadian rhythm and blunt the effects of dawn to sunset fasting. A randomized controlled study conducted on late dinner (11 pm) and routine dinner (6 pm) eaters showed that the postprandial period in late dinner eaters had higher glucose and lower free fatty acids and fatty acid oxidation likely due to altered cortisol rhythm (21).

Healthy sleep patterns are as important as circadian compliant meal timing. Dawn to sunset fasting requires wake-up before dawn to have a pre-dawn breakfast (22). An average of seven or more hours of daily sleep at night is recommended for adults (23). Therefore, individuals who practice dawn to sunset fasting should ensure a minimum of seven hours of sleep at night to prevent sleep deprivation. Taken altogether, practicing dawn to sunset fasting requires self-discipline to perfectly align the mealtimes and wake-up and sleep times with human biological dawn and dusk.

Ramadan fasting (15) is dawn to sunset fasting that is practiced in the holy month of Ramadan (the month of Ramadan is based on the lunar calendar, and therefore it varies every year) for an average of 12-20 hours a day, varying based on geographic location and season for 29 to 30 consecutive days. In contrast to dawn to sunset fasting, the other types of intermittent fasting (e.g., 16:8 intermittent fasting with 16 hours fasting and 8 hours eating) start and end at a self-

determined/predetermined time during the day without taking human biological dawn and dusk into account (24, 25). There are no randomized controlled clinical trials that compared the metabolic effects of dawn to sunset fasting with the effects of other types of intermittent fasting that starts and ends at a self-determined (or predetermined time) without taking biological dawn and dusk into account. A small non-randomized study of time-restricted eating with a fasting period that started and ended at a self-determined time during the day (approximately 14 hours fasting and 10 hours eating) for 12 weeks showed significant improvement in the components of metabolic syndrome (26). In contrast with the findings of this non-randomized study, a large randomized controlled trial conducted in subjects with body mass index between 27 and 43 kg/m<sup>2</sup> showed that time-restricted eating with a fasting period that started and ended at a predetermined time (fasting for 16 hours from 8 pm until 12 pm next day by skipping breakfast and allowing water, tea, coffee during fasting period) for 12 weeks showed no significant changes in the body weight, fat mass, insulin resistance, hemoglobin A1c, and lipids compared with eating three structured meals a day and also snacking between the meals (25).

Another unique feature of dawn to sunset fasting is strict fasting without food or fluid intake during the fasting period and no dietary or calorie restriction during the eating period. In several studies of intermittent fasting other than the dawn to sunset fasting, subjects were reported to perform intermittent fasting or time-restricted eating; however, they did not strictly fast during the fasting period, but instead, they consumed low or no calorie-containing food or drinks (24, 25, 27). In these studies, the effect of intermittent fasting might be limited to caloric restriction rather than the circadian rhythm reset as the subjects were allowed to have low or no calorie-containing food and drinks during the fasting period, likely blunting the effect of mealtime to function as a dominant Zeitgeber (a time giver, rhythmic environmental cue) (28) on peripheral circadian

oscillators. Randomized controlled trials are needed to compare the metabolic effects of dawn to sunset fasting with and without zero-calorie drink intake during the fasting period.

Fasting from dawn to sunset during activity hours may have more potent anti-cancer and immune function enhancing effects than fasting during inactivity or both inactivity and activity hours. Food restriction in mice during the activity period (i.e., 12-hour dark phase in nocturnal animals corresponding to dawn to dusk in humans) was shown to have a more potent anti-cancer effect compared with food restriction during inactivity hours (12-hour light phase) and ad libitum eating (29), suggesting the presence of an optimized immune response associated with fasting during activity hours. The results of intermittent fasting studies conducted on rodents need to be interpreted in the context that rodents are nocturnal animals, and their activity period is nighttime, corresponding to the period from dawn to sunset in humans (17). Two human studies showed that dawn to sunset fasting for 4 weeks induced an anti-cancer proteome in healthy subjects and subjects with metabolic syndrome (16, 17). Randomized, controlled studies are needed to evaluate the effect of dawn to sunset fasting and other types of intermittent fasting on the immune system and cancer prevention.

This review presents scientific literature on the effects of dawn to sunset fasting, a specific type of intermittent fasting commonly practiced during the month of Ramadan, on immune system and how it may mediate physiologic responses against SARS-CoV-2 (**Table 1**).

### **3. EFFECT OF DAWN TO SUNSET FASTING ON IMMUNE SYSTEM**

#### **3.1. Effect of dawn to sunset fasting on inflammation**

SARS-CoV-2 enters host cells via interaction with the angiotensin-converting enzyme 2 (ACE2) protein, markedly expressed in the alveolar epithelium of the lungs and enterocytes of the small intestine (30-32). The immune response to SARS-CoV-2 is similar to other coronaviruses and involves both the innate and adaptive arms of immunity (25). It has been proposed that in mild cases of COVID-19, the inflammatory response generated by the immune system is self-limited as the innate and adaptive immune responses are quick (33). In contrast to mild cases, severe cases of COVID-19 result from delayed and dysregulated innate and adaptive immune responses, which involve recruitment of a massive number of inflammatory cells and generation of high levels of pro-inflammatory cytokines and chemokines, also known as a cytokine storm (33). This impaired immune response leads to severe damage, including acute respiratory distress syndrome, multisystem organ failure, and death (34, 35).

Type I interferon deficiency, mitochondrial dysfunction, and virus-induced senescence were shown to be the driving force of this cytokine storm that leads to severe COVID-19 infection (36-39). Although circulating interferon-beta levels were undetectable in all COVID-19 patients regardless of the severity of the infection, interferon-alpha 2 levels were significantly reduced in patients with critical COVID-19 infection compared with the levels in patients with mild to moderate infection (36). Additionally, low interferon-alpha 2 plasma level was an independent risk factor for progression to critical COVID-19 infection (36).

Persistent elevation of IL-6 and IL-1 levels have been associated with non-survival in COVID-19 (40). Increased IL-6 was shown to be associated with cytokine release syndrome and severe pneumonia (41, 42). These results are in line with the findings of another study that showed that IL-6-induced genes, including IL6R, SOCS3, and STAT3, were significantly expressed in patients stratified as having severe and critical disease than healthy controls (36). A study

conducted in 41 patients with COVID-19 in Wuhan, China, showed that patients admitted to ICU had significantly higher plasma interleukin (IL)-2, IL-7, IL-10, interferon gamma-induced protein 10 (IP-10), which is also known as C-X-C motif chemokine ligand 10 (CXCL10), granulocyte colony-stimulating factor (G-CSF), macrophage inflammatory protein-1A (MIP-1A), monocyte chemoattractant protein-1 (MCP-1), and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) levels compared with patients who were not admitted to ICU (43). A larger study conducted in Wuhan, China, among 452 patients with COVID-19 showed that patients with severe infection had significantly higher IL-2R, IL-6, IL-8, IL-10, and TNF- $\alpha$  than those without severe infection (35).

Several studies of dawn to sunset fasting showed anti-inflammatory effect by suppressing pro-inflammatory cytokines in various populations, including healthy subjects, subjects with at least one component of metabolic syndrome, and subjects with nonalcoholic fatty liver disease (NAFLD) (**Table 1**). A study conducted on 50 healthy subjects who fasted from dawn to sunset for 23 to 30 consecutive days during the month of Ramadan showed that circulating cytokines including IL-6, IL-1 $\beta$ , TNF- $\alpha$  significantly decreased after three weeks of dawn to sunset fasting compared with the baseline levels (44). Likewise, another study of dawn to sunset fasting conducted among healthy subjects showed that plasma levels of cytokines including IL-6, IL-8, IL-1 $\beta$  were significantly reduced after one week of dawn to sunset fasting outside the month of Ramadan and after two weeks of dawn to sunset fasting during the month of Ramadan compared with baseline plasma levels before the weeks of fasting (45). Additionally, the cosinor analysis of cytokines assessing the circadian rhythms of IL-6, IL-8, IL-1 $\beta$  at five different time points during the day (at 11 pm, 2 am, 4 am, 6 am, and 11 am) showed no significant variation in the acrophase of these cytokines during different times of the day (45). The strength of this study was the control of the confounders that could affect the circulating cytokine levels, including sleep duration and

timing, food composition, calories consumed, energy expenditure, and exposure to light (45). More importantly, this study demonstrated that pro-inflammatory cytokines would decrease even after one or two weeks of dawn to sunset fasting (45).

The beneficial effect of dawn to sunset fasting on cytokines also was shown in overweight and obese subjects and subjects with NAFLD. A study that included 82 patients with NAFLD showed that the reduction in serum levels of IL-6 was significantly higher in subjects who fasted from dawn to sunset for a month than the reduction in the IL-6 levels in subjects who did not fast (46). A randomized controlled study conducted in 28 male subjects with obesity showed that 14 subjects who fasted from dawn to sunset for one month had a significantly lower plasma IL-6 level on the 15<sup>th</sup> day of one-month dawn to sunset fasting, 1 and 21 days after the end of one-month dawn to sunset fasting compared with the plasma levels before one-month dawn to sunset fasting (47). In contrast to these findings, there was no significant change in IL-6 levels in control subjects who did not fast (47). A study of 57 overweight and obese subjects who performed dawn to sunset fasting showed a significant reduction in the serum levels of pro-inflammatory cytokines including IL-6, TNF- $\alpha$ , and insulin-like growth factor (IGF-1), along with a significant increase in the anti-inflammatory cytokine IL-10 and IL-10/IL-6 ratio at the end of 23 to 30 days of dawn to sunset fasting compared with the baseline values (48). This study also showed reduction in IGF-1 levels at the end of 23 to 30 days of dawn to sunset fasting compared with the baseline values (48). IGF-1 promotes inflammation in peripheral mononuclear and endothelial cells (49, 50), and several other studies of dawn to sunset fasting reported reduction of IGF-1 with fasting (51, 52).

Regarding the effect of dawn to sunset fasting on chemokines and inflammatory biomarkers, a study of 58 healthy individuals between the ages of 20 to 40 years showed a significant decrease in serum levels of CXCL1, CXCL10, and CXCL12 at the end of one month

dawn to sunset fasting compared with the levels collected on the first day of one-month fasting (53). Although most dawn to sunset fasting studies showed a significant reduction in C-reactive protein levels (46, 54-56), a small study showed no significant change in the C-reactive protein levels (57).

In mice, SARS-CoV-2 causes cytokine storm and lung injury by activating the inflammasome called NOD-, LRR- and pyrin domain-containing protein 3 (NLRP3) through the structural nucleocapsid protein (N protein) (58). The inhibition of NLRP3 inflammasome alleviates cytokine storm and lung injury (58). Two significant mechanisms likely play a role in inhibiting NLRP3 inflammasome during intermittent fasting. The first mechanism appears to be the production of beta-hydroxybutyric acid, a ketone metabolite that was shown to alleviate IL-1 $\beta$  and IL-18 production and caspase-1 activation induced by NLRP3 inflammasome (59). A second mechanism works by activating nuclear receptor subfamily 1 group D member 1 (NR1D1), a core clock component that was shown to inhibit the NLRP3 inflammasome signaling pathway (60). Dawn to sunset fasting for 30 days in humans upregulated the NR1D1, the inhibitor of NLRP3 inflammasome (16). Further human studies of dawn to sunset fasting are needed to replicate these findings in patients with COVID-19.

Altogether, these studies highlight the role of dawn to sunset fasting in promoting an anti-inflammatory milieu by suppressing circulating cytokine, chemokine, pro-inflammatory biomarker levels, and NLRP3 inflammasome. Additionally, the anti-inflammatory effect of dawn to sunset fasting appears to be independent of caloric intake, sleep/wake patterns, and blood collection timing (45). However, larger studies of dawn to sunset fasting need to be conducted controlling for caloric intake and circadian rhythm variations.

### 3.2. Effect of dawn to sunset fasting on oxidative stress

SARS-CoV-2 can lead to mitochondrial dysfunction and the generation of excessive amounts of mitochondrial reactive oxygen species, leading to endothelial dysfunction associated with complications of severe COVID-19 infection (e.g., thrombosis, acute respiratory distress syndrome, and cardiovascular disease) (38, 61, 62). Multiple studies showed that dawn to sunset fasting reduces the generation of reactive oxygen species, which mediate damage to host tissue (52, 63-65). It appears that dawn to sunset fasting reduces oxidative stress by upregulating the expression of antioxidant genes, including transcription factor A, mitochondrial (TFAM), superoxide dismutase 2, mitochondrial (SOD2), and nuclear factor erythroid 2-related factor 2 (Nrf2) (52). A study conducted in 40 subjects with hypertension and 40 controls showed significant improvement in both groups in oxidative stress parameters, including an increase in blood glutathione and decrease malondialdehyde levels at the end of 4-week dawn to sunset fasting and six weeks after completion of 4-week dawn to sunset fasting compared with the levels before 4-week dawn to sunset fasting (63). A similar study conducted in 40 subjects with diabetes and 40 subjects without diabetes showed similar improvement in oxidative stress parameters with 4-week dawn to sunset fasting (64). In contrast to the findings of these two large studies, a small study of dawn to sunset fasting that controlled for sleep patterns and caloric intake and conducted in only eight healthy subjects showed no significant change in malondialdehyde levels and weight (66). These findings could be related to the early measurement of oxidative stress parameters only after two weeks of dawn to sunset fasting instead of 4 weeks of dawn to sunset fasting, inadequate washout time between fasting periods in the month preceding the month of Ramadan and in the month of Ramadan, and lack of weight loss (66) as one study linked alterations in oxidative stress parameters to changes in body weight after fasting (67). Additionally, demonstration of a



substantial difference between before and after Ramadan fasting values of metabolic parameters in healthy subjects can be more difficult compared with subjects with chronic metabolic problems as healthy subjects may have normal or near-normal values of these parameters as the baseline.

### **3.3. Effect of dawn to sunset fasting on autophagy**

Autophagy, a lysosome-dependent cellular process, is crucial for innate and adaptive immunity (68-72). Specifically, autophagy is necessary for the proliferation, differentiation, and function of T cells, B cells, dendritic cells, macrophages, natural killer cells, and generation of chemokines, cytokines, and antibodies (68-72). Downregulation of autophagy is a key mechanism adopted by SARS-CoV-2, to escape detection and have unchecked replication (73). On the other hand, increased autophagy is reported by downregulation of the PRKCSH gene (74) that encodes for hepatocystin (75). Four-week dawn to sunset fasting induced a proteome response associated with increased autophagy simultaneous to reducing oxidative stress and inflammation biomarkers in subjects with metabolic syndrome (17). The reduction in protein kinase C substrate 80K-H (PRKCSH) gene protein products at the end of the 4<sup>th</sup> week during 4-week dawn to sunset fasting and then an average of a 73-fold increase in its level one week after 4-week dawn to sunset fasting suggests increased autophagy during 4-week dawn to sunset fasting, and decreased autophagy with ad libitum eating after 4-week dawn to sunset fasting (17).

### **3.4. Effect of dawn to sunset fasting on immune function**

Studies showed that patients with COVID-19 had increased leukocyte and neutrophil counts along with decreased lymphocyte and monocyte counts (5, 76-79). Additionally, leukocytosis and lymphopenia were more severe in critically ill patients than those with mild infection (3, 5, 76-79). Specifically, lymphopenia was shown to be an independent predictor of

poor prognosis in patients with COVID-19 (42, 79). A study conducted among 452 patients with COVID-19 in Wuhan, China, showed a significant increase in leukocyte numbers, neutrophil to lymphocyte ratio, neutrophil percentage, and a decrease in lymphocyte and monocyte percentages (35). The mean combined number of T, B, and natural killer cells was below the normal range, and this decrease was more profound in severe cases than non-severe cases (35). The mean number of T helper and T suppressor cells was below the normal range in all patients with COVID-19 (35). However, T helper cells were significantly lower in severe cases than in non-severe cases (35). Most importantly, patients with severe COVID-19 infection had a higher naïve to memory T cell ratio than those without severe infection, suggesting that they failed their naïve T cells to differentiate into memory cells (35).

Few studies of dawn to sunset fasting investigated the impact of fasting on peripheral blood cells. A study conducted on 50 healthy subjects who fasted from dawn to sunset for a month had a significant reduction in total leukocytes, granulocytes, lymphocytes, and monocytes after 3 weeks of dawn to sunset fasting compared with baseline levels (44). The same study showed a significant increase in monocyte count and no significant change in other immune cells one month after the completion of the fasting month compared with baseline levels (44). A similar significant increase in monocyte count (55) and no change in leukocyte counts 3 to 4 weeks after completion of one month dawn to sunset fasting month (55, 80). Collectively, the lack of data on the effect of dawn to sunset on the percentage of T cell subsets and T cell function were the significant limitations of these studies. Proteomic analysis of dawn to sunset fasting conducted on subjects with metabolic syndrome showed an average 16-fold increase in the level of calreticulin (CALR) gene protein one week after 4-week dawn to sunset fasting compared with the level before 4-week dawn to sunset fasting (17). Calreticulin (74), a phagocytosis-promoter protein (81) fused with

SARS-CoV S protein 450–650 fragment, was shown to enhance IgG-mediated immune response and result in much higher immunogenicity than SARS-CoV S protein alone (82). Although these findings suggest that dawn to sunset fasting can play an essential role in adaptive humoral defense against SARS-CoV-2 (17), they need to be replicated in SARS-CoV-2.

### **3.5. Effect of dawn to sunset fasting on obesity and metabolic syndrome**

Obesity and metabolic syndrome are major risk factors for severe COVID-19 infection (83-86). Obesity and metabolic syndrome impair adaptive and innate immune responses (87, 88). The excess adipocytes lead to increased pro-inflammatory adipokines, including adiponectin, leptin, IL-6, TNF- $\alpha$ , and IGF-1 (48, 89-91). A study conducted on 417 patients showed that even minor elevation within the normal fasting blood glucose range was a risk factor for admission to the intensive care unit (92). Another study conducted on 941 patients with COVID-19 reported that patients with diabetes had higher mortality compared with those without diabetes, and a fasting blood glucose level equal or greater than 7.0 mmol/L was an independent risk factor for predicting mortality in COVID-19 patients without a history of diabetes (93).

Dawn to sunset fasting improves insulin signaling, blood pressure, and lipid profile with a notable significant increase in high-density lipoprotein levels and leads to an anti-inflammatory effect via a decrease in adipose tissue and weight in the absence of strict regulation of caloric intake or food quality (94-101). Several studies of dawn to sunset fasting conducted in subjects with obesity, type 2 diabetes mellitus, or metabolic syndrome showed that dawn to sunset fasting improves components of metabolic syndrome (17, 47, 57, 98). The improvement in metabolic components was associated with decreased pro-inflammatory cytokines and oxidative stress parameters and upregulation of the genes that play a pivotal role in immune defense (48, 63, 64,

102). A study conducted on 14 subjects with metabolic syndrome showed a significant reduction in body weight, body mass index, and waist circumference at the end of the 4<sup>th</sup> week during 4-week dawn to sunset fasting and one week after 4-week dawn to sunset fasting (17). This study also showed significant improvement in systolic, diastolic, and mean arterial blood pressures at the end of the 4<sup>th</sup> week during 4-week dawn to sunset fasting and a significant reduction in homeostatic model assessment for insulin resistance one week after 4-week dawn to sunset fasting (17). Another study showed that the levels of brain-derived neurotrophic factor (BDNF), a food intake regulator (103) decreased during 4-week dawn to sunset fasting compared with baseline BDNF levels, and there was a positive correlation between changes in the BDNF and TNF- $\alpha$  levels at the end of the 4<sup>th</sup> week during 4-week dawn to sunset fasting suggesting that BDNF may have a role in inflammation attenuated by dawn to sunset fasting (102). As dawn to sunset fasting leads to improvement in metabolic parameters; this is one physiologic way of decreasing the risk of morbidity and mortality associated with COVID-19 (83-86).

### **3.6. Effect of dawn to sunset fasting on the gut microbiome**

A healthy immune system has been linked to a healthy gut microbiome (104). However, patients with COVID-19 were shown to have a significantly altered gut microbiome compared with those without infection (105). In COVID-19, the gut microbiota composition was associated with the disease severity and plasma levels of pro-inflammatory cytokines and chemokines and remained altered even after the disease resolution (105). Other studies showed altered gut microbiome and a positive association between altered gut microbiota and severity of COVID-19 (106-108).

Studies of dawn to sunset fasting showed findings suggestive of beneficial alterations in the gut microbiome (109, 110). A study conducted on both healthy young and middle-aged cohorts showed a significantly increased gut microbiome diversity and remodeling of the gut microbiome in the young cohort during and at the end of dawn to sunset fasting month compared to baseline levels. In the middle-aged cohort, there was a significant remodeling of the gut microbiome and a significant upregulation of Lachnospiraceae and Ruminococcaceae at the end of dawn to sunset fasting month compared with baseline levels. The middle-aged cohort who did not fast had no change in microbiome composition and almost no change in taxa during the study period (109). Another study on 34 healthy subjects showed that dawn to sunset fasting affected beta diversity (assessment of variation in microbial composition between samples) rather than alpha diversity (assessment of diversity and evenness in a sample) (111). While this study found a significant increase in Bacteroidetes and a decrease in Firmicutes in the Pakistani group, a similar effect was not observed in the Chinese group, likely related to the dietary effect on the microbiome (111). A smaller study showed a significant beta diversity and enrichment in the Bacteroidetes phylum after 29 days of dawn to sunset fasting (110). The enrichment in Bacteroidetes phylum after dawn to sunset fasting is important because the reduction in Bacteroidetes/Firmicutes ratio was shown to play an important role in the development of obesity (112, 113). Both Lachnospiraceae and Ruminococcaceae are bacteria that produce butyric acid that play a role in reducing oxidative stress, inflammation, and risk of colon cancer (114-118). Collectively, these studies suggest that dawn to sunset fasting induces favorable changes in the composition of the gut microbiome that can boost the innate and adaptive immune responses. More extensive studies of dawn to sunset fasting are needed to further define the role of the gut microbiome on the immune system controlling for diet and circadian rhythm parameters.

#### **4. CONCERNS REGARDING DAWN TO SUNSET FASTING DURING COVID-19 PANDEMIC**

There is a concern that intermittent fasting leads to dehydration which can affect the mucociliary clearance of respiratory microbial organisms and cause broncho-pulmonary diseases (119). However, no significant data support impaired mucociliary function in dawn to sunset fasting. A study compared the mucociliary clearance time in 40 subjects who fasted for an average of 15 hours from "dawn to sunset" for 29 consecutive days (Ramadan fasting) and 26 subjects who fasted "non-stop" for 60 consecutive hours (Nineveh fasting) (120). In subjects who practiced dawn to sunset Ramadan fasting, there was no significant impairment in mucociliary clearance time four weeks after the Ramadan fasting period compared with time measured during the fasting period (120). In contrast to subjects who practiced dawn to sunset fasting, subjects who practiced Nineveh fasting had a significant prolongation of mucociliary clearance time four weeks after the fasting period compared with the time measured during the fasting period (120).

A second common concern with dawn to sunset fasting is dehydration that can potentially result in renal dysfunction. Several studies of dawn to sunset fasting found no significant dehydration or impairments in renal function in healthy individuals, subjects with metabolic syndrome, as well as in renal transplant patients (121-124). It is important to recognize that during the COVID-19 pandemic, the use of masks and high-grade respirators for prolonged periods, especially in high-risk exposure settings, such as healthcare systems, may cause dehydration from fasting. Dehydration can result from the heat effects of prolonged personal protective equipment (PPE) use and the lack of opportunities to rest and have food and water breaks during the non-fasting period. As a result, dehydration can occur from the high demands of front-line work, which leads to a longer than the recommended period of fasting (125, 126).

A third common concern with dawn to sunset fasting is potential sleep deprivation due to early wake-up time for a pre-dawn breakfast. It is well-known that sleep deprivation can weaken immunity due to dysregulated functioning of immune cells and their responses (127-130). Specifically, nighttime sleep deprivation can disrupt the circadian rhythm and increase oxidative stress (131). Individuals practicing dawn to sunset fasting should have an early bedtime to ensure seven or more hours of daily sleep at night to prevent sleep deprivation.

Individuals with chronic medical conditions (e.g., diabetes, chronic kidney disease, coronary artery disease) should consult their physicians before practicing dawn to sunset fasting/Ramadan fasting and follow medical advice (22, 132, 133). Patients with diabetes mellitus can be at risk for hypoglycemia if they fast long hours and the dose of their medications are not adjusted for mealtimes of dawn to sunset fasting (22, 132, 133). International Diabetes Federation the Ramadan Nutrition Plan (RNP) for patients with diabetes provides a well-balanced nutrition strategy during dawn to sunset fasting/Ramadan fasting that can be considered not only for patients with diabetes but also for individuals without diabetes who plan to practice dawn to sunset fasting (134).

Altogether, during the COVID-19 pandemic, it is crucial to follow safe practices for healthy dawn to sunset fasting/Ramadan fasting (135).

## **5. CONCLUSIONS**

We have summarized the various physiologic effects of dawn to sunset fasting on the immune system. Overall, the health and immune benefits of dawn to sunset fasting not only have the potential to optimize the immune system function during the COVID-19 pandemic, but they can also lead to an improved metabolic profile, decreased risk of cancers, and other chronic

inflammatory diseases. Studies are needed to assess and confirm the potential benefits of dawn to sunset fasting against SARS-CoV-2 directly.

Journal Pre-proof



**Table 1. Representative studies on the impact of dawn to sunset fasting on inflammation, oxidative stress, autophagy, immune function, obesity, metabolic syndrome, components of metabolic syndrome, and gut microbiome**

	Study	Study design	Study population	Duration of dawn to sunset fasting	Summary of study outcome
<b>Inflammation</b>	Faris <i>et al.</i> (2012) (44)	A non-randomized, within-subject design	50 Healthy subjects, 21 men and 29 women, mean (SD) age: 32.7 (9.5) years	Men: 30 days, women: 23 to 25 days during the month of Ramadan	IL-6, IL-1 $\beta$ , TNF- $\alpha$ significantly decreased after 3 weeks of dawn to sunset fasting compared with the baseline levels.
	Almeneesier <i>et al.</i> (2019) (45)	A non-randomized, within-subject design	12 Healthy male subjects, mean (SD) age: 25.1 (2.5) years	1 week outside the month of Ramadan and the first 2 weeks in the month of Ramadan	IL-6, IL-8, IL-1 $\beta$ were significantly reduced after one week of dawn to sunset fasting outside the month of Ramadan and after two weeks of dawn to sunset fasting during the month of Ramadan compared with plasma levels before the weeks of fasting (there was a washout period between dawn to sunset fasting weeks outside the month of Ramadan and during the month of Ramadan).
	Faris <i>et al.</i> (2019) (48)	A non-randomized, within-subject design	57 Subjects with body mass index greater than 25 kg/m <sup>2</sup> , 35 men and 22 women, mean (SD) age: 36.2 (12.5) years	Men: 28 to 30 days, women: 23 to 25 days during the month of Ramadan	There was a significant reduction in the serum levels of pro-inflammatory cytokines, including IL-6, TNF- $\alpha$ , and insulin-like growth factor (IGF-1), along with a significant increase in the anti-inflammatory cytokine IL-10 and IL-10/IL-6 ratio at the end of 23 to 30 days of dawn to sunset fasting compared with the baseline values.
	Rahbar <i>et al.</i> (2019) (51)	A non-randomized, within-subject design	34 Male subjects were excluded if they were smokers, took thyroid and hypertension medications as well as estrogens, had clinical dyslipidemia, diabetes mellitus, and hypothyroidism, 34 men, mean age (SD): 35 (11) years	One month during the month of Ramadan	There was a significant decrease in circulating IGF-1 and IL-2 levels at the end of one month of dawn to sunset fasting compared with the levels before one-month dawn to sunset fasting.

Madkour <i>et al.</i> (2019) (52)	A non-randomized, within-subject design	56 Subjects with body mass index greater than 25 kg/m <sup>2</sup> , 34 men and 22 women, controls (n=6) with healthy body weight, mean (SD) age: 35.72 (12.35) years in overweight/obese subjects, 29.8 (14) years in controls	One month during the month of Ramadan	There was a significant reduction in IGF-1 levels at the end of one month of dawn to sunset fasting compared with the levels before one month dawn to sunset fasting.
Aliasghari <i>et al.</i> (2017) (46)	Non-randomized, controlled, within-subject, and between-subject design	83 Subjects with nonalcoholic fatty liver disease (NAFLD), 42 subjects who fasted, 41 subjects who did not, 57 men and 26 women, mean (SD) age: 37.59 (7.06) years in fasting group, 35.80 (7.33) years in non-fasting group	One month during the month of Ramadan	The reduction in serum levels of IL-6 and high-sensitivity C-reactive protein was significantly higher in subjects who fasted from dawn to sunset for a month than the reduction in the levels of these pro-inflammatory markers in subjects who did not fast
Akrami Mohajeri <i>et al.</i> (2013) (53)	A non-randomized, within-subject design	58 Healthy male subjects, age: 20 to 40 years	One month during the month of Ramadan	There was a significant decrease in serum levels of CXCL1, CXCL10, and CXCL12 at the end of one month of dawn to sunset fasting compared with the levels collected on the first day of one-month fasting.
Mari <i>et al.</i> (2021) (54)	A non-randomized, controlled, within-subject design/retrospective, case-control study	74 subjects with NAFLD who fasted and 81 subjects who did not fast, mean age (SD): 51.8 (20.9) years in fasting group, 52.6 (19.3) years in non-fasting group	One month during the month of Ramadan	There was a significant reduction in C-reactive protein levels after dawn to sunset fasting compared with the levels before dawn to sunset fasting.
Zouhal <i>et al.</i> (2020) (47)	A randomized controlled, within-subject, and between-subject design	28 Male subjects, 14 in the fasting group, 14 in the non-fasting group, with obesity with body mass index between 30 and 40 kg/m <sup>2</sup> , mean (SD) age: 24 (3.4) years in the intervention group, 23.8 (3.8) years in the control group	One month during the month of Ramadan	Subjects who fasted from dawn to sunset for one month had a significantly lower plasma IL-6 level on the 15th day of one-month dawn to sunset fasting, 1 day and 21 days after the end of one-month dawn to sunset fasting compared with the plasma levels before one-month dawn to sunset fasting. In contrast to these findings, there was no

					significant change in IL-6 levels in control subjects who did not fast.
	Mindikoglu <i>et al.</i> (2020) (16)	A non-randomized, within-subject design	14 Healthy subjects, 13 men and 1 woman, mean age: 32 years	Men: 30 days, woman: 26 days during the month of Ramadan	Dawn to sunset fasting upregulated the nuclear receptor subfamily 1 group D member 1 (NR1D1) which is the inhibitor of NLRP3 inflammasome. There was an average 11 fold increase in the NR1D1 protein level at the end of 4th week of 30-day dawn to sunset fasting compared with the level before 30-day dawn to sunset fasting.
<b>Oxidative stress</b>	Madkour <i>et al.</i> (2019) (52)	A non-randomized, within-subject design	56 Subjects with body mass index greater than 25 kg/m <sup>2</sup> , controls with healthy body weight, 34 men and 22 women, 6 controls, mean (SD) age: 35.72 (12.35) years in overweight/obese subjects, 29.8 (14) years in controls	One month during the month of Ramadan	There was a significant increase in the relative expression of the antioxidant genes, including transcription factor A, mitochondrial (TFAM), superoxide dismutase 2, mitochondrial (SOD2), and nuclear factor erythroid 2-related factor 2 (Nrf2) compared with the expression of the antioxidant genes in the controls at the end of one month of dawn to sunset fasting in subjects who fasted.
	Al-Shafei (2014) (63)	A non-randomized, controlled, within-subject design	40 Subjects with hypertension and 40 controls, mean age (SD): 55 (5), both men and women in equal proportions	One month during the month of Ramadan	In both groups, there was a significant improvement in both groups in oxidative stress parameters, including an increase in blood glutathione and decrease malondialdehyde levels during the 4th week of dawn to sunset fasting and six weeks after completion of 4-week dawn to sunset fasting compared with the levels before 4-week dawn to sunset fasting.
	Al-Shafei (2014) (64)	A non-randomized, controlled, within-subject design	40 Subjects with diabetes and 40 without diabetes, mean age (SD): 55 (5), both men and women in equal proportions	One month during the month of Ramadan	In both groups, there was a significant improvement in both groups in oxidative stress parameters, including an increase in blood glutathione and decrease malondialdehyde levels during the 4th week of dawn to sunset fasting and six weeks after

					completion of 4-week dawn to sunset fasting compared with the levels before 4-week dawn to sunset fasting.
<b>Autophagy</b>	Mindikoglu <i>et al.</i> (2020) (17)	A non-randomized, within-subject design	14 subjects with metabolic syndrome, 8 men and 6 women, mean age (SD): 59 years (16)	29 days during the month of Ramadan	Four-week dawn to sunset fasting induced a proteome response associated with increased autophagy simultaneous to the reduction in oxidative stress and inflammation biomarkers in subjects with metabolic syndrome. The reduction in protein kinase C substrate 80K-H (PRKCSH) gene protein products at the end of the 4th week during 4-week dawn to sunset fasting and then an average of a 73-fold increase in its level one week after 4-week dawn to sunset fasting suggests increased autophagy during 4-week dawn to sunset fasting, and decreased autophagy with ad libitum eating after 4-week dawn to sunset fasting.
<b>Immune function</b>	Faris <i>et al.</i> (2012) (44)	A non-randomized, within-subject design	50 Healthy subjects, 21 men and 29 women, mean (SD) age: 32.7 (9.5) years	Men: 30 days, women: 23 to 25 days during the month of Ramadan	There was a significant reduction in total leukocytes, granulocytes, lymphocytes, and monocytes after 3 weeks of dawn to sunset fasting compared with baseline levels. The same study showed a significant increase in monocyte count and no significant change in other immune cells one month after the completion of the fasting month compared with baseline levels.
	Mindikoglu <i>et al.</i> (2020) (17)	A non-randomized, within-subject design	14 subjects with metabolic syndrome, 8 men and 6 women, mean age (SD): 59 years (16)	29 days during the month of Ramadan	There was an average 16-fold increase in the level of calreticulin (CALR) gene protein one week after 4-week dawn to sunset fasting compared with the level before 4-week dawn to sunset fasting.
<b>Obesity, metabolic syndrome, components of metabolic syndrome</b>	Mindikoglu <i>et al.</i> (2020) (17)	A non-randomized, within-subject design	14 subjects with metabolic syndrome, 8 men and 6 women, mean age (SD): 59 years (16)	29 days during the month of Ramadan	There was a significant reduction in body weight, body mass index, waist circumference both at the end of the 4th week during 4-week dawn to sunset fasting and one week after 4-week dawn to sunset fasting. This study also showed significant improvement in systolic, diastolic, and mean

					arterial blood pressures at the end of the 4th week during 4-week dawn to sunset fasting and a significant reduction in homeostatic model assessment for insulin resistance one week after 4-week dawn to sunset fasting.
Faris <i>et al.</i> (2019) (48)	A non-randomized, within-subject design	57 Subjects with body mass index greater than 25 kg/m <sup>2</sup> , 35 men and 22 women, mean (SD) age: 36.2 (12.5) years	Men: 28 to 30 days, women: 23 to 25 days during the month of Ramadan		There was a significant reduction in total cholesterol, triacylglycerol, visceral fat surface area, body mass index, fat mass, body weight, and systolic blood pressure after 3 weeks of dawn to sunset fasting compared with the baseline levels.
Al-Shafei (2014) (63)	A non-randomized, controlled, within-subject design	40 Subjects with hypertension and 40 controls, mean age (SD): 55 (5), both men and women in equal proportions	One month during the month of Ramadan		In subjects with hypertension, there was a significant reduction in systolic and diastolic blood pressure, triglyceride, and low-density lipoprotein levels and a significant increase in high-density lipoprotein levels during the 4th week of dawn to sunset fasting compared with the levels before 4-week dawn to sunset fasting. In control subjects, there was a significant decrease in triglyceride levels during the 4th week of dawn to sunset fasting compared with the levels before 4-week dawn to sunset fasting.
Al-Shafei (2014) (64)	A non-randomized, controlled, within-subject design	40 Subjects with diabetes and 40 without diabetes, mean age (SD): 55 (5), both men and women in equal proportions	One month during the month of Ramadan		In subjects with diabetes, there was a significant improvement in fasting blood glucose and triglyceride levels during the 4th week of dawn to sunset fasting compared with the levels before 4-week dawn to sunset fasting. In subjects without diabetes, there was a significant decrease in triglyceride levels during the 4th week of dawn to sunset fasting compared with the levels before 4-week dawn to sunset fasting.
Tahapary <i>et al.</i> (2020) (98)	Meta-analysis (28 observational studies)	Subjects with type 2 diabetes mellitus, age (SD): 48.67 to 60.1	10 to 29-30 days during the month of Ramadan		There was a significant decrease in fasting glucose level (11 studies), hemoglobin A1c (16 studies), decrease in total cholesterol, low-density lipoprotein, high-density lipoprotein, and triglyceride levels (11 studies), body weight (15 studies), and waist

					circumference (6 studies) at the end of dawn to sunset fasting period compared with baseline levels.
	Nematy <i>et al.</i> (2012) (101)	A non-randomized, within-subject design	82 subjects with coronary artery disease, metabolic syndrome, or cerebrovascular disease, 38 men and 44 women, age (SD): 54 (10)	One month during the month of Ramadan	There was a significant reduction in body weight, body mass index, waist circumference, total cholesterol, triglycerides, low-density lipoprotein, systolic blood pressure, and a significant increase in high-density lipoprotein levels at the end of dawn to sunset fasting (from the 27th day of dawn to sunset to 6 days after the end of the fasting month) compared with the levels from before the fasting month (from the 7 days before the start of fasting month until 2 first days of the fasting month).
<b>Gut microbiome</b>	Su <i>et al.</i> (2021) (109)	A non-randomized, controlled, within-subject design	30 healthy young male adult cohort who fasted, age (SD): 18.63 (1.75), 27 healthy middle-aged cohort who fasted, age (SD): 39.9 (6.4), 10 healthy middle-aged cohort who did not fast, age (SD): 42.6 (7.9)	One month during the month of Ramadan	In the young cohort, there was a significantly increased gut microbiome diversity and remodeling of the gut microbiome during and at the end of dawn to sunset fasting month compared to baseline levels. In the middle-aged cohort, there was a significant remodeling of the gut microbiome and a significant upregulation of Lachnospiraceae and Ruminococcaceae at the end of dawn to sunset fasting month compared with baseline levels. The middle-aged cohort who did not fast had no change in microbiome composition and almost no change in taxa during the study period.
	Ozkul <i>et al.</i> (2020) (110)	A non-randomized, within-subject design	9 healthy subjects, 2 men and 7 women, age (SD): 45 (9.7)	29 days during the month of Ramadan	There was a significant beta diversity and enrichment in the Bacteroidetes phylum after 29 days of dawn to sunset fasting.
	Ali <i>et al.</i> (2021) (111)	A non-randomized, within-subject design	34 healthy subjects (16 Chinese and 18 Pakistani), age range 18 to 40 years	One month during the month of Ramadan	There was a significant increase in Bacteroidetes and a decrease in Firmicutes in the Pakistani group, and a significant increase in Proteobacteria and a decrease in Bacteroidetes in the Chinese cohort after

					down to sunset fasting month compared to the levels before fasting month.
--	--	--	--	--	---

Journal Pre-proof

## **CONFLICT OF INTEREST**

None of the authors has a conflict of interest.

Journal Pre-proof



## REFERENCES

1. WHO Coronavirus (COVID-19) Dashboard. Available at <https://covid19.who.int/>. Accessed on December 22, 2021.
2. Booth A, Reed AB, Ponzo S, Yassaee A, Aral M, Plans D, et al. Population risk factors for severe disease and mortality in COVID-19: A global systematic review and meta-analysis. *PloS one*. 2021;16(3):e0247461.
3. Gao YD, Ding M, Dong X, Zhang JJ, Kursat Azkur A, Azkur D, et al. Risk factors for severe and critically ill COVID-19 patients: A review. *Allergy*. 2021;76(2):428-55.
4. Melo AKG, Milby KM, Caparroz ALMA, Pinto ACPN, Santos RRP, Rocha AP, et al. Biomarkers of cytokine storm as red flags for severe and fatal COVID-19 cases: A living systematic review and meta-analysis. *PloS one*. 2021;16(6):e0253894.
5. Zhang J-j, Cao Y-y, Tan G, Dong X, Wang B-c, Lin J, et al. Clinical, radiological, and laboratory characteristics and risk factors for severity and mortality of 289 hospitalized COVID-19 patients. *Allergy*. 2021;76(2):533-50.
6. van Eijk LE, Binkhorst M, Bourgonje AR, Offringa AK, Mulder DJ, Bos EM, et al. COVID-19: immunopathology, pathophysiological mechanisms, and treatment options. *The Journal of Pathology*. 2021;254(4):307-31.
7. Rubin EJ, Longo DL. SARS-CoV-2 Vaccination — An Ounce (Actually, Much Less) of Prevention. *New England Journal of Medicine*. 2020;383(27):2677-8.
8. Wang Z, Fu Y, Guo Z, Li J, Li J, Cheng H, et al. Transmission and prevention of SARS-CoV-2. *Biochemical Society transactions*. 2020;48(5):2307-16.

9. Faris MAE, Salem ML, Jahrami HA, Madkour MI, BaHammam AS. Ramadan intermittent fasting and immunity: An important topic in the era of COVID-19. *Annals of thoracic medicine*. 2020;15(3):125-33.
10. Javanmard SH, Otroj Z. Ramadan Fasting and Risk of Covid-19. *International journal of preventive medicine*. 2020;11:60.
11. Ealey KN, Phillips J, Sung H-K. COVID-19 and obesity: fighting two pandemics with intermittent fasting. *Trends in Endocrinology & Metabolism*. 2021;32(9):706-20.
12. Gnoni M, Beas R, Vázquez-Garagatti R. Is there any role of intermittent fasting in the prevention and improving clinical outcomes of COVID-19?: intersection between inflammation, mTOR pathway, autophagy and calorie restriction. *VirusDisease*. 2021.
13. Moghadam MT, Taati B, Paydar Ardakani SM, Suzuki K. Ramadan Fasting During the COVID-19 Pandemic; Observance of Health, Nutrition and Exercise Criteria for Improving the Immune System. *Frontiers in Nutrition*. 2021;7(349).
14. Hannan MA, Rahman MA, Rahman MS, Sohag AAM, Dash R, Hossain KS, et al. Intermittent fasting, a possible priming tool for host defense against SARS-CoV-2 infection: Crosstalk among calorie restriction, autophagy and immune response. *Immunology letters*. 2020;226:38-45.
15. Fasting times. QUR'AN. Surah Al-Baqarah (2:187).
16. Mindikoglu AL, Abdulsada MM, Jain A, Choi JM, Jalal PK, Devaraj S, et al. Intermittent fasting from dawn to sunset for 30 consecutive days is associated with anticancer proteomic signature and upregulates key regulatory proteins of glucose and lipid metabolism, circadian clock, DNA repair, cytoskeleton remodeling, immune system and cognitive function in healthy subjects. *J Proteomics*. 2020;217:103645.

17. Mindikoglu AL, Abdulsada MM, Jain A, Jalal PK, Devaraj S, Wilhelm ZR, et al. Intermittent fasting from dawn to sunset for four consecutive weeks induces anticancer serum proteome response and improves metabolic syndrome. *Sci Rep.* 2020;10(1):18341.
18. Wehr TA, Aeschbach D, Duncan WC, Jr. Evidence for a biological dawn and dusk in the human circadian timing system. *The Journal of physiology.* 2001;535(Pt 3):937-51.
19. Wehr TA, Schwartz PJ, Turner EH, Feldman-Naim S, Drake CL, Rosenthal NE. Bimodal patterns of human melatonin secretion consistent with a two-oscillator model of regulation. *Neuroscience letters.* 1995;194(1-2):105-8.
20. Witbracht M, Keim NL, Forester S, Widaman A, Laugero K. Female breakfast skippers display a disrupted cortisol rhythm and elevated blood pressure. *Physiol Behav.* 2015;140:215-21.
21. Gu C, Brereton N, Schweitzer A, Cotter M, Duan D, Børsheim E, et al. Metabolic Effects of Late Dinner in Healthy Volunteers-A Randomized Crossover Clinical Trial. *The Journal of clinical endocrinology and metabolism.* 2020;105(8):2789-802.
22. Mindikoglu AL, Opekun AR, Gagan SK, Devaraj S. Impact of Time-Restricted Feeding and Dawn-to-Sunset Fasting on Circadian Rhythm, Obesity, Metabolic Syndrome, and Nonalcoholic Fatty Liver Disease. *Gastroenterology research and practice.* 2017;2017:3932491.
23. Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, Buysse D, et al. Recommended Amount of Sleep for a Healthy Adult: A Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society. *Sleep.* 2015;38(6):843-4.
24. Kunduraci YE, Ozbek H. Does the Energy Restriction Intermittent Fasting Diet Alleviate Metabolic Syndrome Biomarkers? A Randomized Controlled Trial. *Nutrients.* 2020;12(10).

25. Lowe DA, Wu N, Rohdin-Bibby L, Moore AH, Kelly N, Liu YE, et al. Effects of Time-Restricted Eating on Weight Loss and Other Metabolic Parameters in Women and Men With Overweight and Obesity: The TREAT Randomized Clinical Trial. *JAMA Internal Medicine*. 2020;180(11):1491-9.
26. Wilkinson MJ, Manoogian ENC, Zadourian A, Lo H, Fakhouri S, Shoghi A, et al. Ten-Hour Time-Restricted Eating Reduces Weight, Blood Pressure, and Atherogenic Lipids in Patients with Metabolic Syndrome. *Cell Metab*. 2020;31(1):92-104.e5.
27. Harney DJ, Hutchison AT, Hatchwell L, Humphrey SJ, James DE, Hocking S, et al. Proteomic Analysis of Human Plasma during Intermittent Fasting. *Journal of proteome research*. 2019;18(5):2228-40.
28. Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev*. 2000;14(23):2950-61.
29. Wu MW, Li XM, Xian LJ, Levi F. Effects of meal timing on tumor progression in mice. *Life Sci*. 2004;75(10):1181-93.
30. Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol*. 2004;203(2):631-7.
31. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, et al. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature*. 2003;426(6965):450-4.
32. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-3.

33. Sokolowska M, Lukasik ZM, Agache I, Akdis CA, Akdis D, Akdis M, et al. Immunology of COVID-19: Mechanisms, clinical outcome, diagnostics, and perspectives-A report of the European Academy of Allergy and Clinical Immunology (EAACI). *Allergy*. 2020;75(10):2445-76.
34. Channappanavar R, Perlman S. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Seminars in immunopathology*. 2017;39(5):529-39.
35. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, et al. Dysregulation of Immune Response in Patients With Coronavirus 2019 (COVID-19) in Wuhan, China. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2020;71(15):762-8.
36. Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Smith N, et al. Impaired type I interferon activity and inflammatory responses in severe COVID-19 patients. *Science*. 2020;369(6504):718-24.
37. Lee S, Yu Y, Trimpert J, Benthani F, Mairhofer M, Richter-Pechanska P, et al. Virus-induced senescence is a driver and therapeutic target in COVID-19. *Nature*. 2021.
38. Chang R, Mamun A, Dominic A, Le N-T. SARS-CoV-2 Mediated Endothelial Dysfunction: The Potential Role of Chronic Oxidative Stress. *Frontiers in Physiology*. 2021;11(1752).
39. Blanco-Melo D, Nilsson-Payant BE, Liu WC, Uhl S, Hoagland D, Møller R, et al. Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. *Cell*. 2020;181(5):1036-45.e9.

40. McGonagle D, Sharif K, O'Regan A, Bridgewood C. The Role of Cytokines including Interleukin-6 in COVID-19 induced Pneumonia and Macrophage Activation Syndrome-Like Disease. *Autoimmunity reviews*. 2020;19(6):102537.
41. Wang W, Liu X, Wu S, Chen S, Li Y, Nong L, et al. Definition and Risks of Cytokine Release Syndrome in 11 Critically Ill COVID-19 Patients With Pneumonia: Analysis of Disease Characteristics. *The Journal of infectious diseases*. 2020;222(9):1444-51.
42. Zhang X, Tan Y, Ling Y, Lu G, Liu F, Yi Z, et al. Viral and host factors related to the clinical outcome of COVID-19. *Nature*. 2020;583(7816):437-40.
43. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet (London, England)*. 2020;395(10223):497-506.
44. Faris MA, Kacimi S, Al-Kurd RA, Fararjeh MA, Bustanji YK, Mohammad MK, et al. Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects. *Nutrition research (New York, NY)*. 2012;32(12):947-55.
45. Almeneessier AS, BaHammam AA, Alzoghaibi M, Olaish AH, Nashwan SZ, BaHammam AS. The effects of diurnal intermittent fasting on proinflammatory cytokine levels while controlling for sleep/wake pattern, meal composition and energy expenditure. *PloS one*. 2019;14(12):e0226034.
46. Aliasghari F, Izadi A, Gargari BP, Ebrahimi S. The Effects of Ramadan Fasting on Body Composition, Blood Pressure, Glucose Metabolism, and Markers of Inflammation in NAFLD Patients: An Observational Trial. *Journal of the American College of Nutrition*. 2017;36(8):640-5.

47. Zouhal H, Bagheri R, Ashtary-Larky D, Wong A, Triki R, Hackney AC, et al. Effects of Ramadan intermittent fasting on inflammatory and biochemical biomarkers in males with obesity. *Physiology & Behavior*. 2020;225:113090.
48. Faris MAE, Madkour MI, Obaideen AK, Dalah EZ, Hasan HA, Radwan H, et al. Effect of Ramadan diurnal fasting on visceral adiposity and serum adipokines in overweight and obese individuals. *Diabetes research and clinical practice*. 2019;153:166-75.
49. Che W, Lerner-Marmarosh N, Huang Q, Osawa M, Ohta S, Yoshizumi M, et al. Insulin-Like Growth Factor-1 Enhances Inflammatory Responses in Endothelial Cells. *Circulation research*. 2002;90(11):1222-30.
50. Wolters TLC, Netea MG, Hermus A, Smit JWA, Netea-Maier RT. IGF1 potentiates the pro-inflammatory response in human peripheral blood mononuclear cells via MAPK. *Journal of molecular endocrinology*. 2017;59(2):129-39.
51. Rahbar AR, Safavi E, Rooholamini M, Jaafari F, Darvishi S, Rahbar A. Effects of Intermittent Fasting during Ramadan on Insulin-like Growth Factor-1, Interleukin 2, and Lipid Profile in Healthy Muslims. *International journal of preventive medicine*. 2019;10:7.
52. Madkour MI, A TE-S, Jahrami HA, Sherif NM, Hassan RE, Awadallah S, et al. Ramadan diurnal intermittent fasting modulates SOD2, TFAM, Nrf2, and sirtuins (SIRT1, SIRT3) gene expressions in subjects with overweight and obesity. *Diabetes research and clinical practice*. 2019;155:107801.
53. Akrami Mohajeri F, Ahmadi Z, Hassanshahi G, Akrami Mohajeri E, Ravari A, Ghalebi SR. Dose Ramadan Fasting Affects Inflammatory Responses: Evidences for Modulatory Roles of This Unique Nutritional Status via Chemokine Network. *Iranian journal of basic medical sciences*. 2013;16(12):1217-22.

54. Mari A, Khoury T, Baker M, Said Ahmad H, Abu Baker F, Mahamid M. The Impact of Ramadan Fasting on Fatty Liver Disease Severity: A Retrospective Case Control Study from Israel. *The Israel Medical Association journal* : IMAJ. 2021;23(2):94-8.
55. Askari VR, Alavinezhad A, Boskabady MH. The impact of "Ramadan fasting period" on total and differential white blood cells, haematological indices, inflammatory biomarker, respiratory symptoms and pulmonary function tests of healthy and asthmatic patients. *Allergologia et immunopathologia*. 2016;44(4):359-67.
56. Aksungar FB, Topkaya AE, Akyildiz M. Interleukin-6, C-reactive protein and biochemical parameters during prolonged intermittent fasting. *Annals of nutrition & metabolism*. 2007;51(1):88-95.
57. Sadiya A, Ahmed S, Siddieg HH, Babas IJ, Carlsson M. Effect of Ramadan fasting on metabolic markers, body composition, and dietary intake in Emiratis of Ajman (UAE) with metabolic syndrome. *Diabetes, metabolic syndrome and obesity : targets and therapy*. 2011;4:409-16.
58. Pan P, Shen M, Yu Z, Ge W, Chen K, Tian M, et al. SARS-CoV-2 N protein promotes NLRP3 inflammasome activation to induce hyperinflammation. *Nature communications*. 2021;12(1):4664.
59. Youm YH, Nguyen KY, Grant RW, Goldberg EL, Bodogai M, Kim D, et al. The ketone metabolite  $\beta$ -hydroxybutyrate blocks NLRP3 inflammasome-mediated inflammatory disease. *Nature medicine*. 2015;21(3):263-9.
60. Pourcet B, Zecchin M, Ferri L, Beauchamp J, Sitaula S, Billon C, et al. Nuclear Receptor Subfamily 1 Group D Member 1 Regulates Circadian Activity of NLRP3 Inflammasome to



Reduce the Severity of Fulminant Hepatitis in Mice. *Gastroenterology*. 2018;154(5):1449-64.e20.

61. Mehri F, Rahbar AH, Ghane ET, Souri B, Esfahani M. The comparison of oxidative markers between Covid-19 patients and healthy subjects. *Archives of medical research*. 2021.
62. Pincemail J, Cavalier E, Charlier C, Cheramy-Bien JP, Brevers E, Courtois A, et al. Oxidative Stress Status in COVID-19 Patients Hospitalized in Intensive Care Unit for Severe Pneumonia. A Pilot Study. *Antioxidants (Basel, Switzerland)*. 2021;10(2).
63. Al-Shafei AI. Ramadan fasting ameliorates arterial pulse pressure and lipid profile, and alleviates oxidative stress in hypertensive patients. *Blood pressure*. 2014;23(3):160-7.
64. Al-Shafei AI. Ramadan fasting ameliorates oxidative stress and improves glyceimic control and lipid profile in diabetic patients. *European journal of nutrition*. 2014;53(7):1475-81.
65. Faris MeA-IE, Jahrami HA, Obaideen AA, Madkour MI. Impact of diurnal intermittent fasting during Ramadan on inflammatory and oxidative stress markers in healthy people: Systematic review and meta-analysis. *Journal of Nutrition & Intermediary Metabolism*. 2019;15:18-26.
66. BaHammam AS, Pandi-Perumal SR, Alzoghaibi MA. The effect of Ramadan intermittent fasting on lipid peroxidation in healthy young men while controlling for diet and sleep: A pilot study. *Annals of thoracic medicine*. 2016;11(1):43-8.
67. Faris MA, Hussein RN, Al-Kurd RA, Al-Fararjeh MA, Bustanji YK, Mohammad MK. Impact of ramadan intermittent fasting on oxidative stress measured by urinary 15-f(2t)-isoprostane. *Journal of nutrition and metabolism*. 2012;2012:802924.
68. Pan H, Chen L, Xu Y, Han W, Lou F, Fei W, et al. Autophagy-associated immune responses and cancer immunotherapy. *Oncotarget*. 2016;7(16):21235-46.

69. Crotzer VL, Blum JS. Autophagy and adaptive immunity. *Immunology*. 2010;131(1):9-17.
70. Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. *Nature reviews Molecular cell biology*. 2018;19(6):349-64.
71. Cui B, Lin H, Yu J, Yu J, Hu Z. Autophagy and the Immune Response. *Advances in experimental medicine and biology*. 2019;1206:595-634.
72. Jiang GM, Tan Y, Wang H, Peng L, Chen HT, Meng XJ, et al. The relationship between autophagy and the immune system and its applications for tumor immunotherapy. *Molecular cancer*. 2019;18(1):17.
73. Gassen NC, Papies J, Bajaj T, Emanuel J, Dethloff F, Chua RL, et al. SARS-CoV-2-mediated dysregulation of metabolism and autophagy uncovers host-targeting antivirals. *Nature communications*. 2021;12(1):3818.
74. Gene [Internet]. Bethesda (MD): National Library of Medicine (US), National Center for Biotechnology Information; 2004 – [cited 2021 Dec 04]. Available from: <https://www.ncbi.nlm.nih.gov/gene/>.
75. Yang J, Zhao Y, Ma K, Jiang FJ, Liao W, Zhang P, et al. Deficiency of hepatocystin induces autophagy through an mTOR-dependent pathway. *Autophagy*. 2011;7(7):748-59.
76. Lee J, Park SS, Kim TY, Lee DG, Kim DW. Lymphopenia as a Biological Predictor of Outcomes in COVID-19 Patients: A Nationwide Cohort Study. *Cancers*. 2021;13(3).
77. Mazzoni A, Salvati L, Maggi L, Annunziato F, Cosmi L. Hallmarks of immune response in COVID-19: Exploring dysregulation and exhaustion. *Seminars in immunology*. 2021:101508.
78. Huang G, Kovalic AJ, Graber CJ. Prognostic Value of Leukocytosis and Lymphopenia for Coronavirus Disease Severity. *Emerging infectious diseases*. 2020;26(8):1839-41.

79. Tan L, Wang Q, Zhang D, Ding J, Huang Q, Tang Y-Q, et al. Lymphopenia predicts disease severity of COVID-19: a descriptive and predictive study. *Signal Transduction and Targeted Therapy*. 2020;5(1):33.
80. Maughan RJ, Leiper JB, Bartagi Z, Zrifi R, Zerguini Y, Dvorak J. Effect of Ramadan fasting on some biochemical and haematological parameters in Tunisian youth soccer players undertaking their usual training and competition schedule. *Journal of sports sciences*. 2008;26 Suppl 3:S39-46.
81. Raghavan M, Wijeyesakere SJ, Peters LR, Del Cid N. Calreticulin in the immune system: ins and outs. *Trends in immunology*. 2013;34(1):13-21.
82. Qiu X, Hong C, Li Y, Bao W, Gao X-M. Calreticulin as a hydrophilic chimeric molecular adjuvant enhances IgG responses to the spike protein of severe acute respiratory syndrome coronavirus. *Microbiology and Immunology*. 2012;56(8):554-61.
83. Gao F, Zheng KI, Wang X-B, Sun Q-F, Pan K-H, Wang T-Y, et al. Obesity Is a Risk Factor for Greater COVID-19 Severity. *Diabetes Care*. 2020;43(7):e72-e4.
84. Steenblock C, Schwarz PEH, Ludwig B, Linkermann A, Zimmet P, Kulebyakin K, et al. COVID-19 and metabolic disease: mechanisms and clinical management. *The lancet Diabetes & endocrinology*. 2021;9(11):786-98.
85. Xu Y, Yang X, Bian H, Xia M. Metabolic dysfunction associated fatty liver disease and coronavirus disease 2019: clinical relationship and current management. *Lipids in health and disease*. 2021;20(1):126.
86. Zhang J, Kong W, Xia P, Xu Y, Li L, Li Q, et al. Impaired Fasting Glucose and Diabetes Are Related to Higher Risks of Complications and Mortality Among Patients With Coronavirus Disease 2019. *Frontiers in endocrinology*. 2020;11:525.

87. Andersen CJ, Murphy KE, Fernandez ML. Impact of Obesity and Metabolic Syndrome on Immunity. *Advances in nutrition* (Bethesda, Md). 2016;7(1):66-75.
88. Wang Q, Wu H. T Cells in Adipose Tissue: Critical Players in Immunometabolism. *Frontiers in immunology*. 2018;9:2509.
89. Tilg H, Moschen AR. Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Nature reviews Immunology*. 2006;6(10):772-83.
90. Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose Expression of Tumor Necrosis Factor- $\alpha$ ; Direct Role in Obesity-Linked Insulin Resistance. *Science*. 1993;259(5091):87-91.
91. Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM. Increased adipose tissue expression of tumor necrosis factor- $\alpha$  in human obesity and insulin resistance. *The Journal of Clinical Investigation*. 1995;95(5):2409-15.
92. Alahmad B, Al-Shammari AA, Bennakhi A, Al-Mulla F, Ali H. Fasting Blood Glucose and COVID-19 Severity: Nonlinearity Matters. *Diabetes Care*. 2020;43(12):3113-6.
93. Cai Y, Shi S, Yang F, Yi B, Chen X, Li J, et al. Fasting blood glucose level is a predictor of mortality in patients with COVID-19 independent of diabetes history. *Diabetes research and clinical practice*. 2020;169:108437.
94. Fernando HA, Zibellini J, Harris RA, Seimon RV, Sainsbury A. Effect of Ramadan Fasting on Weight and Body Composition in Healthy Non-Athlete Adults: A Systematic Review and Meta-Analysis. *Nutrients*. 2019;11(2).
95. Jahrami HA, Alsibai J, Clark CCT, Faris MAE. A systematic review, meta-analysis, and meta-regression of the impact of diurnal intermittent fasting during Ramadan on body weight in healthy subjects aged 16 years and above. *European journal of nutrition*. 2020;59(6):2291-316.

96. Kul S, Savaş E, Öztürk ZA, Karadağ G. Does Ramadan fasting alter body weight and blood lipids and fasting blood glucose in a healthy population? A meta-analysis. *Journal of religion and health*. 2014;53(3):929-42.
97. Sadeghirad B, Motaghipisheh S, Kolahdooz F, Zahedi MJ, Haghdoost AA. Islamic fasting and weight loss: a systematic review and meta-analysis. *Public health nutrition*. 2014;17(2):396-406.
98. Tahapary DL, Astrella C, Kristanti M, Harbuwono DS, Soewondo P. The impact of Ramadan fasting on metabolic profile among type 2 diabetes mellitus patients: A meta-analysis. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2020;14(5):1559-70.
99. Adlouni A, Ghalim N, Benslimane A, Lecerf JM, Saile R. Fasting during Ramadan induces a marked increase in high-density lipoprotein cholesterol and decrease in low-density lipoprotein cholesterol. *Annals of nutrition & metabolism*. 1997;41(4):242-9.
100. Adlouni A, Ghalim N, Saile R, Hda N, Parra HJ, Benslimane A. Beneficial effect on serum apo AI, apo B and Lp AI levels of Ramadan fasting. *Clinica chimica acta; international journal of clinical chemistry*. 1998;271(2):179-89.
101. Nematy M, Alinezhad-Namaghi M, Rashed MM, Mozhdehifard M, Sajjadi SS, Akhlaghi S, et al. Effects of Ramadan fasting on cardiovascular risk factors: a prospective observational study. *Nutr J*. 2012;11:69.
102. Abdulsada MM, Wilhelm ZR, Opekun AR, Devaraj S, Jalal PK, Mindikoglu AL. The effect of four-week intermittent fasting from dawn to sunset on circulating brain-derived neurotrophic factor levels in subjects with metabolic syndrome and healthy subjects. *Metabolism Open*. 2021;9:100070.

103. Rosas-Vargas H, Martínez-Ezquerro JD, Bienvenu T. Brain-derived neurotrophic factor, food intake regulation, and obesity. *Archives of medical research*. 2011;42(6):482-94.
104. Wiertsema SP, van Bergenhenegouwen J, Garssen J, Knippels LMJ. The Interplay between the Gut Microbiome and the Immune System in the Context of Infectious Diseases throughout Life and the Role of Nutrition in Optimizing Treatment Strategies. *Nutrients*. 2021;13(3).
105. Yeoh YK, Zuo T, Lui GC, Zhang F, Liu Q, Li AY, et al. Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19. *Gut*. 2021;70(4):698-706.
106. Yamamoto S, Saito M, Tamura A, Prawisuda D, Mizutani T, Yotsuyanagi H. The human microbiome and COVID-19: A systematic review. *PloS one*. 2021;16(6):e0253293.
107. Zuo T, Zhang F, Lui GCY, Yeoh YK, Li AYL, Zhan H, et al. Alterations in Gut Microbiota of Patients With COVID-19 During Time of Hospitalization. *Gastroenterology*. 2020;159(3):944-55.e8.
108. Gu S, Chen Y, Wu Z, Chen Y, Gao H, Lv L, et al. Alterations of the Gut Microbiota in Patients With Coronavirus Disease 2019 or H1N1 Influenza. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2020;71(10):2669-78.
109. Su J, Wang Y, Zhang X, Ma M, Xie Z, Pan Q, et al. Remodeling of the gut microbiome during Ramadan-associated intermittent fasting. *The American journal of clinical nutrition*. 2021;113(5):1332-42.
110. Ozkul C, Yalinay M, Karakan T. Structural changes in gut microbiome after Ramadan fasting: a pilot study. *Beneficial microbes*. 2020;11(3):227-33.

111. Ali I, Liu K, Long D, Faisal S, Hilal MG, Ali I, et al. Ramadan Fasting Leads to Shifts in Human Gut Microbiota Structured by Dietary Composition. *Frontiers in Microbiology*. 2021;12(314).
112. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature*. 2006;444(7122):1027-31.
113. Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI. Obesity alters gut microbial ecology. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;102(31):11070-5.
114. Hamer HM, Jonkers DM, Bast A, Vanhoutvin SA, Fischer MA, Kodde A, et al. Butyrate modulates oxidative stress in the colonic mucosa of healthy humans. *Clinical nutrition (Edinburgh, Scotland)*. 2009;28(1):88-93.
115. Hamer HM, Jonkers D, Venema K, Vanhoutvin S, Troost FJ, Brummer RJ. Review article: the role of butyrate on colonic function. *Alimentary pharmacology & therapeutics*. 2008;27(2):104-19.
116. Scharlau D, Borowicki A, Habermann N, Hofmann T, Klenow S, Miene C, et al. Mechanisms of primary cancer prevention by butyrate and other products formed during gut flora-mediated fermentation of dietary fibre. *Mutation research*. 2009;682(1):39-53.
117. Bui TP, Ritari J, Boeren S, de Waard P, Plugge CM, de Vos WM. Production of butyrate from lysine and the Amadori product fructoselysine by a human gut commensal. *Nature communications*. 2015;6:10062.
118. Rajilić-Stojanović M, de Vos WM. The first 1000 cultured species of the human gastrointestinal microbiota. *FEMS microbiology reviews*. 2014;38(5):996-1047.

119. Kalhoff H. Mild dehydration: a risk factor of broncho-pulmonary disorders? *European Journal of Clinical Nutrition*. 2003;57(2):S81-S7.
120. Develioglu ON, Sirazi S, Topak M, Purisa S, Kulekci M. Differences in Mucociliary activity of volunteers undergoing Ramadan versus Nineveh fasting. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*. 2013;270(5):1655-9.
121. Abdalla AH, Shaheen FA, Rassoul Z, Owda AK, Popovich WF, Mousa DH, et al. Effect of Ramadan fasting on Moslem kidney transplant recipients. *American journal of nephrology*. 1998;18(2):101-4.
122. Cheah SH, Ch'ng SL, Husain R, Duncan MT. Effects of fasting during Ramadan on urinary excretion in Malaysian Muslims. *The British journal of nutrition*. 1990;63(2):329-37.
123. Leiper JB, Molla AM, Molla AM. Effects on health of fluid restriction during fasting in Ramadan. *Eur J Clin Nutr*. 2003;57 Suppl 2:S30-8.
124. Aslam M, Wilson JV. Clinical problems during the fast of Ramadan. *Lancet (London, England)*. 1989;1(8644):955.
125. Manjunath G, Aravindhakshan R, Varghese S. Effect of fasting during Ramadan on thermal stress parameters. *Eastern Mediterranean health journal = La revue de sante de la Mediterranee orientale = al-Majallah al-sihhiyah li-sharq al-mutawassit*. 2019;25(1):34-9.
126. Schmahl FW, Metzler B. The health risks of occupational stress in islamic industrial workers during the Ramadan fasting period. *Polish journal of occupational medicine and environmental health*. 1991;4(3):219-28.



127. Ajabnoor GMA, Bahijri S, Shaik NA, Borai A, Alamoudi AA, Al-Aama JY, et al. Ramadan fasting in Saudi Arabia is associated with altered expression of CLOCK, DUSP and IL-1alpha genes, as well as changes in cardiometabolic risk factors. PloS one. 2017;12(4):e0174342.
128. Bryant PA, Trinder J, Curtis N. Sick and tired: Does sleep have a vital role in the immune system? Nature reviews Immunology. 2004;4(6):457-67.
129. Wilder-Smith A, Mustafa FB, Earnest A, Gen L, Macary PA. Impact of partial sleep deprivation on immune markers. Sleep medicine. 2013;14(10):1031-4.
130. Yousfi N, Bragazzi NL, Briki W, Zmijewski P, Chamari K. The COVID-19 pandemic: how to maintain a healthy immune system during the lockdown - a multidisciplinary approach with special focus on athletes. Biology of sport. 2020;37(3):211-6.
131. Teixeira KRC, dos Santos CP, de Medeiros LA, Mendes JA, Cunha TM, De Angelis K, et al. Night workers have lower levels of antioxidant defenses and higher levels of oxidative stress damage when compared to day workers. Scientific Reports. 2019;9(1):4455.
132. International Diabetes Federation Diabetes and Ramadan Practical Guidelines 2021. Available at <https://www.idf.org/our-activities/education/diabetes-and-ramadan/healthcare-professionals.html>. Accessed on December 9, 2021.
- .
133. Ibrahim M, Davies MJ, Ahmad E, Annabi FA, Eckel RH, Ba-Essa EM, et al. Recommendations for management of diabetes during Ramadan: update 2020, applying the principles of the ADA/EASD consensus. BMJ Open Diabetes Research & Care. 2020;8(1):e001248.

134. International Diabetes Federation Diabetes and Ramadan Practical Guidelines 2021.

Chapter 8. The Ramadan Nutrition Plan (RNP) for people with diabetes. Hamdy O, Yusof BNM, Maher S.

[https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwiWy7SLhdj0AhXOk2oFHbF0CuYQFnoECAgQAQ&url=https%3A%2F%2Fwww.idf.org%2Fcomponent%2Fattachments%2F%3Ftask%3Ddownload%26id%3D2351%3A210126\\_DaR\\_CH8\\_The-Ramadan-Nutrition-Plan-RNP-for-people-with-diabetes\\_SingleLow&usg=AOvVaw1Ff52jABYKMeiwcZiVr9l](https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwiWy7SLhdj0AhXOk2oFHbF0CuYQFnoECAgQAQ&url=https%3A%2F%2Fwww.idf.org%2Fcomponent%2Fattachments%2F%3Ftask%3Ddownload%26id%3D2351%3A210126_DaR_CH8_The-Ramadan-Nutrition-Plan-RNP-for-people-with-diabetes_SingleLow&usg=AOvVaw1Ff52jABYKMeiwcZiVr9l). Accessed on December 9, 2021.

135. World Health O. World Health Organization. (2021). Safe Ramadan practices in the context of COVID-19: interim guidance, 7 April 2021. World Health Organization.

<https://apps.who.int/iris/handle/10665/340661>. License: CC BY-NC-SA 3.0 IGO. Available at <https://apps.who.int/iris/bitstream/handle/10665/340661/WHO-2019-nCoV-Ramadan-2021.1-eng.pdf?sequence=1&isAllowed=y>. Accessed on December 09, 2021. Geneva: World Health Organization; 2021 2021. Contract No.: WHO/2019-nCoV/Ramadan/2021.1.